8223086

## **ALHCA**



Test Booklet Code



This Booklet contains 24 pages.

Do not open this Test Booklet until you are asked to do so.

Read carefully the Instructions on the Back Cover of this Test Booklet.

## Important Instructions:

- The Answer Sheet is inside this Test Booklet. When you are directed to open the Test Booklet, take out the Answer Sheet and fill in the particulars on Side-1 and Side-2 carefully with blue/black ball point pen only.
- 2. The test is of 3 hours duration and this Test Booklet contains 180 questions. Each question carries 4 marks. For each correct response, the candidate will get 4 marks. For each incorrect response, one mark will be deducted from the total scores. The maximum marks are 720.
- 3. Use Blue/Black Ball Point Pen only for writing particulars on this page/marking responses.
- 4. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
- On completion of the test, the candidate must hand over the Answer Sheet to the Invigilator before leaving the Room/Hall. The candidates are allowed to take away this Test Booklet with them.
- 6. The CODE for this Booklet is **XX**. Make sure that the CODE printed on **Side-2** of the Answer Sheet is the same as that on this Test Booklet. In case of discrepancy, the candidate should immediately report the matter to the Invigilator for replacement of both the Test Booklet and the Answer Sheet.
- 7. The candidates should ensure that the Answer Sheet is not folded. Do not make any stray marks on the Answer Sheet. Do not write your Roll No. anywhere else except in the specified space in the Test Booklet/Answer Sheet.
- 8. Use of white fluid for correction is not permissible on the Answer Sheet.

Name of the Candidate (in Capitals): M. Hohameel Ashoray Ali MOHANED MU	STAFA.
Roll Number: in figures 511904426.	
: in words fifty one croser wentere laters four thousand po	cis openins depostantia
Centre of Examination (in Capitals): TRINITY INTERNATIONAL SCHOOL	(CBSE) 10
Candidate's Signature : M. TROFIDITE Albard Rle Invigilator's Signature :	
Facsimile signature stamp of	
Centre Superintendent :	v
ALHCA/XX/Page 1	English

- The bond dissociation energies of X<sub>2</sub>, Y<sub>2</sub> and XY are in the ratio of 1:0.5:1. ΔH for the formation of XY is -200 kJ mol<sup>-1</sup>. The bond dissociation energy of X<sub>2</sub> will be
  - (1)  $400 \text{ kJ mol}^{-1}$
  - (2) 200 kJ mol<sup>-1</sup>
  - (3) 800 kJ mol<sup>-1</sup>
  - (4) 100 kJ mol<sup>-1</sup>
- When initial concentration of the reactant is doubled, the half-life period of a zero order reaction
  - (1) remains unchanged
  - (2) is halved
  - (3) is tripled
  - (4) is doubled
- 3. The correction factor 'a' to the ideal gas equation corresponds to
  - (1) forces of attraction between the gas molecules
  - (2) density of the gas molecules
  - (3) electric field present between the gas molecules
  - (4) volume of the gas molecules
- 4. Which one of the following conditions will favour maximum formation of the product in the reaction,

$$A_2(g) + B_2(g) \rightleftharpoons X_2(g) \quad \Delta_r H = -X kJ$$
?

- (1) High temperature and low pressure
- (2) Low temperature and high pressure
- (3) High temperature and high pressure
- (4) Low temperature and low pressure
- 5. For the redox reaction

$$MnO_4^- + C_2O_4^{2-} + H^+ \longrightarrow Mn^{2+} + CO_2 + H_2O$$

the correct coefficients of the reactants for the balanced equation are

	$MnO_4^-$	$C_2O_4^{2-}$	H <sup>+</sup>
(1)	5	16	2
(2)	16	5	2
(3)	2	16	5
(4)	2	5	16

6. In the reaction

the electrophile involved is

- (1) dichlorocarbene (:CCl<sub>2</sub>)
- (2) dichloromethyl cation (CHCl₂)
- (3) dichloromethyl anion (CHCl<sub>2</sub>)
- (4) formyl cation (CHO)
- Carboxylic acids have higher boiling points than aldehydes, ketones and even alcohols of comparable molecular mass. It is due to their
  - (1) formation of intermolecular H-bonding
  - (2) formation of intramolecular H-bonding
  - (3) more extensive association of carboxylic acid via van der Waals force of attraction
  - (4) formation of carboxylate ion
- NaOI (produced by reacting Y with NaOH) and yields a yellow precipitate with characteristic smell.

A and Y are respectively

(2) 
$$H_3C$$
  $\sim$   $CH_2$   $OH$  and  $I_2$ 

(3) 
$$CH - CH_3$$
 and  $I_2$  OH

(4) 
$$\sim$$
 CH<sub>2</sub> – CH<sub>2</sub> – OH and I<sub>2</sub>

- 9. The correct difference between first- and second-order reactions is that
  - (1) the rate of a first-order reaction does depend on reactant concentrations; the rate of a second-order reaction does not depend on reactant concentrations
  - (2) the rate of a first-order reaction does not depend on reactant concentrations; the rate of a second-order reaction does depend on reactant concentrations
  - a first-order reaction can be catalyzed; a second-order reaction cannot be catalyzed
  - (4) the half-life of a first-order reaction does not depend on [A]<sub>0</sub>; the half-life of a second-order reaction does depend on [A]<sub>0</sub>
- 10. Consider the change in oxidation state of Bromine corresponding to different emf values as shown in the diagram below:

$$BrO_4^- \xrightarrow{1.82 \text{ V}} BrO_3^- \xrightarrow{1.5 \text{ V}} HBrO$$

$$Br^- \xleftarrow{1.0652 \text{ V}} Br_2 \xleftarrow{1.595 \text{ V}}$$

Then the species undergoing disproportionation is

- (1) HBrO
- (2) BrO $_3^-$
- (3) Br<sub>2</sub>
- (4) BrO<sub>4</sub>
- Among CaH<sub>2</sub>, BeH<sub>2</sub>, BaH<sub>2</sub>, the order of ionic character is
  - (1)  $BaH_2 < BeH_2 < CaH_2$
  - (2)  $BeH_2 < CaH_2 < BaH_2$
  - (3)  $BeH_2 < BaH_2 < CaH_2$
  - (4)  $CaH_2 < BeH_2 < BaH_2$
- 12. In which case is the number of molecules of water maximum?
  - (1)  $10^{-3}$  mol of water
  - (2) 18 mL of water
  - (3) 0.00224 L of water vapours at 1 atm and 273 K
  - (4) 0.18 g of water

- 13. The difference between amylose and amylopectin
  - (1) Amylose is made up of glucose and galactose
  - (2) Amylopectin have  $1 \rightarrow 4$   $\alpha$ -linkage and  $1 \rightarrow 6$   $\alpha$ -linkage
  - (3) Amylopectin have 1  $\rightarrow$  4  $\alpha$ -linkage and 1  $\rightarrow$  6  $\beta$ -linkage
  - (4) Amylose have  $1 \rightarrow 4$   $\alpha$ -linkage and  $1 \rightarrow 6$   $\beta$ -linkage
- Nitration of aniline in strong acidic medium also gives m-nitroaniline because
  - In acidic (strong) medium aniline is present as anilinium ion.
  - (2) In spite of substituents nitro group always goes to only m-position.
  - (3) In absence of substituents nitro group always goes to m-position.
  - (4) In electrophilic substitution reactions amino group is meta directive.
- 15. Which of the following oxides is most acidic in nature?
  - (1) CaO
  - (2) MgO
  - (3) BaO
  - (4) BeO
- 16. A mixture of 2·3 g formic acid and 4·5 g oxalic acid is treated with conc. H<sub>2</sub>SO<sub>4</sub>. The evolved gaseous mixture is passed through KOH pellets. Weight (in g) of the remaining product at STP will be
  - (1) 4.4
  - (2) 1.4
  - (3) 2.8
  - (4) 3.0
- 17. Regarding cross-linked or network polymers, which of the following statements is incorrect?
  - They contain strong covalent bonds in their polymer chains.
  - (2) They contain covalent bonds between various linear polymer chains.
  - (3) Examples are bakelite and melamine.
  - (4) They are formed from bi- and tri-functional monomers.

- 18. Magnesium reacts with an element (X) to form an ionic compound. If the ground state electronic configuration of (X) is 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>3</sup>, the simplest formula for this compound is
  - (1)  $Mg_3X_2$
  - (2)  $Mg_2X_3$
  - (3) Mg<sub>2</sub>X
  - (4) MgX<sub>2</sub>
- 19. Iron exhibits bcc structure at room temperature. Above 900°C, it transforms to fcc structure. The ratio of density of iron at room temperature to that at 900°C (assuming molar mass and atomic radii of iron remains constant with temperature) is
  - (1)  $\frac{1}{2}$
  - $(2) \quad \frac{\sqrt{3}}{\sqrt{2}}$
  - $(3) \quad \frac{3\sqrt{3}}{4\sqrt{2}}$
  - $(4) \quad \frac{4\sqrt{3}}{3\sqrt{2}}$
- 20. Which one is a wrong statement?
  - (1) The value of m for d<sub>2</sub> is zero.
  - (2) Total orbital angular momentum of electron in 's' orbital is equal to zero.
  - (3) The electronic configuration of N atom is

$1s^2$	$2s^2$	$2p_x^1$	$2p_y^1$	$2p_z^1$
↑↓	$\uparrow \downarrow$	1	1	1

- (4) An orbital is designated by three quantum numbers while an electron in an atom is designated by four quantum numbers.
- 21. Consider the following species:

Which one of these will have the highest bond order?

- (1) CN
- (2) NO
- (3) CN+
- (4) CN

- 22. Following solutions were prepared by mixing different volumes of NaOH and HCl of different concentrations:
  - a.  $60 \text{ mL } \frac{\text{M}}{10} \text{ HCl} + 40 \text{ mL } \frac{\text{M}}{10} \text{ NaOH}$
  - b.  $55 \text{ mL } \frac{\text{M}}{10} \text{ HCl} + 45 \text{ mL } \frac{\text{M}}{10} \text{ NaOH}$
  - c.  $75 \text{ mL } \frac{\text{M}}{5} \text{ HCl} + 25 \text{ mL } \frac{\text{M}}{5} \text{ NaOH}$
  - d.  $100 \text{ mL } \frac{M}{10} \text{ HCl} + 100 \text{ mL } \frac{M}{10} \text{ NaOH}$

pH of which one of them will be equal to 1?

- (1)
- (2) b
- (3) d
- (4) a
- **23.** The solubility of  ${\rm BaSO_4}$  in water is  $2\cdot 42\times 10^{-3}~{\rm gL^{-1}}$  at 298 K. The value of its solubility product  $({\rm K_{sp}})$  will be

(Given molar mass of  $BaSO_4 = 233 \text{ g mol}^{-1}$ )

- (1)  $1.08 \times 10^{-8} \text{ mol}^2 \text{ L}^{-2}$
- (2)  $1.08 \times 10^{-10} \text{ mol}^2 \text{ L}^{-2}$
- (3)  $1.08 \times 10^{-14} \text{ mol}^2 \text{ L}^{-2}$
- (4)  $1.08 \times 10^{-12} \text{ mol}^2 \text{ L}^{-2}$
- 24. On which of the following properties does the coagulating power of an ion depend?
  - (1) The sign of charge on the ion alone
  - (2) The magnitude of the charge on the ion alone
  - (3) Both magnitude and sign of the charge on the ion
  - (4) Size of the ion alone
- 25. Given van der Waals constant for NH<sub>3</sub>, H<sub>2</sub>, O<sub>2</sub> and CO<sub>2</sub> are respectively 4·17, 0·244, 1·36 and 3·59, which one of the following gases is most easily liquefied?
  - (1) CO<sub>2</sub>
  - (2) NH<sub>3</sub>
  - (3)  $O_2$
  - (4) H<sub>2</sub>

- 26. The geometry and magnetic behaviour of the complex [Ni(CO)<sub>4</sub>] are
  - (1) tetrahedral geometry and paramagnetic
  - (2) square planar geometry and diamagnetic
  - (3) square planar geometry and paramagnetic
  - (4) tetrahedral geometry and diamagnetic
- 27. Iron carbonyl, Fe(CO)5 is
  - (1) dinuclear
  - (2) tetranuclear
  - (3) trinuclear
  - (4) mononuclear
- 28. Match the metal ions given in Column I with the spin magnetic moments of the ions given in Column II and assign the correct code:

	Colu	mnI		Column II
a.	Co <sup>3+</sup>		i	$\sqrt{8}$ B.M.
b.	Cr3+		ii.	$\sqrt{35}$ B.M.
c.	Fe <sup>3+</sup>		iii.	$\sqrt{3}$ B.M.
d.	Ni <sup>2+</sup>		iv.	$\sqrt{24}$ B.M.
			v.	$\sqrt{15}$ B.M.
	a	b	c	d
(1)	iii	v	i	ii
(2)	iv	v	ii	i
(3)	iv	i	ii	iii
(4)	i	ii	iii	iv

- 29. The type of isomerism shown by the complex [CoCl<sub>2</sub>(en)<sub>2</sub>] is
  - (1) Linkage isomerism
  - (2) Geometrical isomerism
  - (3) Ionization isomerism
  - (4) Coordination isomerism
- 30. Which one of the following ions exhibits d-d transition and paramagnetism as well?
  - (1)  $MnO_4^{2-}$
  - (2)  $CrO_4^{2-}$
  - (3) MnO<sub>4</sub>
  - (4)  $Cr_2O_7^{2-}$

- 31. The compound A on treatment with Na gives B, and with PCl<sub>5</sub> gives C. B and C react together to give diethyl ether. A, B and C are in the order
  - (1) C<sub>2</sub>H<sub>5</sub>OH, C<sub>2</sub>H<sub>5</sub>ONa, C<sub>2</sub>H<sub>5</sub>Cl
  - (2) C<sub>2</sub>H<sub>5</sub>OH, C<sub>2</sub>H<sub>6</sub>, C<sub>2</sub>H<sub>5</sub>Cl
  - (3) C<sub>2</sub>H<sub>5</sub>Cl, C<sub>2</sub>H<sub>6</sub>, C<sub>2</sub>H<sub>5</sub>OH
  - (4) C<sub>2</sub>H<sub>5</sub>OH, C<sub>2</sub>H<sub>5</sub>Cl, C<sub>2</sub>H<sub>5</sub>ONa
- 32. The compound C<sub>7</sub>H<sub>8</sub> undergoes the following reactions:

$$C_7H_8 \xrightarrow{3 \text{ Cl}_2/\Delta} A \xrightarrow{Br_2/\text{ Fe}} B \xrightarrow{Zn/\text{ HCl}} C$$

The product 'C' is

- (1) p-bromotoluene
- (2) m-bromotoluene
- (3) 3-bromo-2,4,6-trichlorotoluene
- (4) o-bromotoluene
- 33. Hydrocarbon (A) reacts with bromine by substitution to form an alkyl bromide which by Wurtz reaction is converted to gaseous hydrocarbon containing less than four carbon atoms. (A) is
  - (1) CH<sub>4</sub>
  - (2) CH = CH
  - (3)  $CH_3 CH_3$
  - (4)  $CH_2 = CH_2$
- 34. Which oxide of nitrogen is **not** a common pollutant introduced into the atmosphere both due to natural and human activity?
  - (1) NO
  - (2)  $N_2O_5$
  - (3) N<sub>2</sub>O
  - (4) NO<sub>2</sub>

35. Identify the major products P, Q and R in the following sequence of reactions:

$$\begin{array}{c} \text{Anhydrous} \\ + \text{CH}_3\text{CH}_2\text{CH}_2\text{Cl} & \xrightarrow{\text{AlCl}_3} \\ \\ P \xrightarrow{\text{(i) O}_2} & \text{Q} + \text{R} \end{array}$$

P Q R

- (1)  $CH(CH_3)_2$   $CH_3 CO CH_2$
- (2)  $CH_2CH_2CH_3$  CHO  $CH_3CH_2 OH$
- (3)  $CH(CH_3)_2$  OH  $CH_3CH(OH)CH_3$

$$(4) \begin{picture}(4){c} $\operatorname{CH}_2\operatorname{CH}_3$ & $\operatorname{CHO}$ & $\operatorname{COOH}$ \\ & & & & \\ \end{picture}$$

- 36. Which of the following compounds can form a zwitterion?
  - (1) Glycine
  - (2) Aniline
  - (3) Benzoic acid
  - (4) Acetanilide

- 37. Which of the following molecules represents the order of hybridisation sp<sup>2</sup>, sp<sup>2</sup>, sp, sp from left to right atoms?
  - (1)  $CH_3 CH = CH CH_3$
  - (2)  $HC \equiv C C \equiv CH$
  - (3)  $CH_2 = CH CH = CH_2$
  - (4)  $CH_2 = CH C = CH$
- 38. Which of the following carbocations is expected to be most stable?

$$(1) \qquad \begin{matrix} H \\ Y \end{matrix} \qquad \begin{matrix} NO_2 \\ \end{matrix}$$

- Which of the following is correct with respect to
   I effect of the substituents? (R = alkyl)
  - $(1) NR_2 > OR > F$
  - (2)  $-NH_2 < -OR < -F$
  - (3)  $-NH_2 > -OR > -F$
  - $(4) NR_2 < -OR < -F$

40.	Which of the following statements is <b>not</b> true for halogens?	46.	rep	resent	s the	lung o	condi		ptions correctl in asthma an
	(1) Chlorine has the highest electron-gain enthalpy.	1	(1)	Dec	na, res reased	resp	oirat		surface;
	(2) All form monobasic oxyacids.			The second	ammati	State Est	-		
	(3) All but fluorine show positive oxidation states.	1	(2)	resp	iratory			onch	ioles; Decrease
	(4) All are oxidizing agents.		(3)		eased ammati	resp on of b			surface;
41.	Considering Ellingham diagram, which of the following metals can be used to reduce alumina?		(4)		eased i iratory			brone	chioles; Increase
	(1) Cu	47.	Mat	ch the	e items	given	in C	olun	n I with those i
	(2) Fe				II and	select	the	corr	rect option give
	(3) Mg		belo		MARKET T			~	1 11
	(4) Zn			50/12	ımn I	340		- 5	olumn II
40	The correct order of atomic radii in group 13		a.	Tric	uspid v	aive	i.		tween left atriun I left ventricle
42.	elements is		b.	Bica	spid va	lve	ii.	Later Control	tween right
	(1) B < Ga < Ál < In < Tl	57		Dice	DPIG 1				tricle and
	(2) B < Al < In < Ga < Tl							-	monary artery
	(3) B < Ga < Al < Tl < In		c.	Sem	ilunar	valve	iii.		tween right
	(4) B < Al < Ga < In < Tl								ium and right atricle
10	NA 90 1000 1 000			a	b	c			
43.	In the structure of ClF <sub>3</sub> , the number of lone pairs of electrons on central atom 'Cl' is	1	(1)	ii	i	iii			
	(1) three		(2)	iii	i	ii			
	(2) one		(3)	i	ii	iii			
	(3) four		(4)	i	iii	ii			
	(4) two	48.	Mat	ch th	e items	given	in (	Colun	nn I with those i
44.	The correct order of N-compounds in it decreasing order of oxidation states is	14.500.000	Col	umn w:.	II and				rect option give
	(1) NH <sub>4</sub> Cl, N <sub>2</sub> , NO, HNO <sub>3</sub>			Coli	ımn I				Column II
	(2) HNO <sub>3</sub> , NO, N <sub>2</sub> , NH <sub>4</sub> Cl		a.	Tida	l volur	ne		i.	2500 – 3000 mI
	(3) HNO <sub>3</sub> , NH <sub>4</sub> Cl, NO, N <sub>2</sub>		b.	Insp	iratory me	Reser	ve	ii.	1100 – 1200 mI
	(4) HNO <sub>3</sub> , NO, NH <sub>4</sub> Cl, N <sub>2</sub>		c.		iratory	Reser	ve	iii.	500 - 550  mL
45.	Which one of the following elements is unable to		(30)	volu		1		2.2	1000 – 1100 mI
	form MF <sub>6</sub> <sup>3-</sup> ion?		d.	a	dual vo	c		d	1000 – 1100 mi
	(1) In		(1)	iv	iii	ii		i	
	(2) Ga		(2)	iii	ii	i		iv	
	(3) B		(3)	i	iv	ii		iii	
	(4) Al	- 81	(4)	iii	i	iv		ii	

49. The transparent lens in the human eye is held in 53. its place by smooth muscles attached to the ciliary body (2) ligaments attached to the ciliary body smooth muscles attached to the iris **50.** 51.

	(4) liga	aments attached	to the iris		(3) (4)		UAUG(				
•	Which o		s an amino acid derived	54.		mn II				umn I with those in orrect option given	
	(1) Est	triol			belov	v : Colui	mn T			Column II	
	(2) Ep	inephrine			250			e Phase		Breakdown of	
		tradiol			a.	Prom	ierative	e rnase	1.	endometrial lining	
	(4) Ecc	lysone			b.	Secre	tory Pl	hase	ii.	Follicular Phase	
					C.	Mens	truatio	on	iii.	Luteal Phase	
		f the following ctly paired with	structures or regions is its function?		( - N	a	b	c			
	(1) Con	rpus callosum	: band of fibers connecting left and right cerebral hemispheres.		(1) (2) (3)	iii iii ii	i ii iii	ii i i			
	(2) Me	dulla oblongata	: controls respiration and cardiovascular reflexes.	55.	(4) All or (1)		iii ollowin moter	ii g are pa	art o	f an operon except	
	(3) Hy	pothalamus	production of releasing hormones and regulation of temperature,		(2) (3) (4)	an en	erator hancer tural g	=			
	(4) Lin	nbic system	hunger and thirst.  : consists of fibre tracts that interconnect different regions of brain; controls movement.	56.		ition i Mino Multi	s r muta iple ste otypic v		ions	, the mechanism o	f
		of the following	hormones can play a orosis?	57.			nas an l			ndition on one of her romosome can be	

AGGTATCGCAT is a sequence from the coding

strand of a gene. What will be the corresponding

sequence of the transcribed mRNA?

UCCAUAGCGUA

AGGUAUCGCAU

(2)

(2)

(3)

Aldosterone and Prolactin

(4) Progesterone and Aldosterone

Parathyroid hormone and Prolactin

Estrogen and Parathyroid hormone

52.

inherited by

Both sons and daughters

Only daughters

Only sons

Only grandchildren

- In which disease does mosquito transmitted 64. pathogen cause chronic inflammation lymphatic vessels? Amoebiasis (1) Elephantiasis

  - Ringworm disease (3)
  - Ascariasis (4)
- Among the following sets of examples for divergent evolution, select the incorrect option :
  - Eye of octopus, bat and man
  - Forelimbs of man, bat and cheetah
  - Brain of bat, man and cheetah
  - Heart of bat, man and cheetah
- Conversion of milk to curd improves its nutritional value by increasing the amount of
  - Vitamin E
  - (2) Vitamin D
  - Vitamin B<sub>12</sub>
  - Vitamin A
- Which of the following is not an autoimmune disease?
  - (1) Vitiligo
  - **Psoriasis**
  - Alzheimer's disease
  - Rheumatoid arthritis
- The similarity of bone structure in the forelimbs of many vertebrates is an example of
  - Adaptive radiation
  - Homology (2)
  - Convergent evolution (3)
  - Analogy
- Which of the following characteristics represent 'Inheritance of blood groups' in humans?
  - **Dominance** a.
  - Co-dominance b.
  - Multiple allele c.
  - Incomplete dominance d.
  - Polygenic inheritance e.
  - (1) a. c and e
  - b, c and e
  - b, d and e
  - a, b and c

Match the items given in Column I with those in Column II and select the correct option given below:

Column I

Column II

- a Eutrophication
- i. UV-B radiation
- Sanitary landfill . ii. Deforestation

  - iii. Nutrient Snow blindness

enrichment

- iv. Waste disposal .Ihum cultivation
  - b d
- (1)
- iv iii (2)
- ii i (3) iii iii
- iv All of the following are included in Ex-situ

ii

Seed banks (1)

conservation' except

(4)

- (2) Wildlife safari parks
- Botanical gardens (3)
- Sacred groves
- one of the following population interactions is widely used in medical science for the production of antibiotics?
  - Amensalism
  - Commensalism
  - Parasitism (3)
  - Mutualism
- In a growing population of a country,
  - pre-reproductive individuals are less than the reproductive individuals.
  - pre-reproductive individuals are more than the reproductive individuals.
  - pre-reproductive reproductive and individuals are equal in number.
  - reproductive individuals are less than the post-reproductive individuals.
- Which part of poppy plant is used to obtain the 68. drug "Smack"?
  - Leaves (1)
  - (2)Flowers
  - Roots
  - (4) Latex

69.	Hormones secreted by the placenta to maintain pregnancy are	73.		ımn I			n Column I with those in the <i>correct</i> option given
	(1) hCG, progestogens, estrogens, glucocorticoids		pero		$mn\ I$		Column II
	(2) hCG, hPL, progestogens, prolactin		a.	Glyc	osuria	i.	Accumulation of uric acid in joints
	<ul><li>(3) hCG, hPL, progestogens, estrogens</li><li>(4) hCG, hPL, estrogens, relaxin, oxytocin</li></ul>	10	b.	Gou		ii.	Mass of crystallised salts within the kidney
70.	The amnion of mammalian embryo is derived		c.	Rena	al calculi	iii.	Inflammation in glomeruli
	from (1) ectoderm and endoderm		d.	100000000000000000000000000000000000000	nerular pritis	iv.	Presence of glucose in urine
	(2) ectoderm and mesoderm			a	b	c	d
	(3) mesoderm and trophoblast		(1)	iv	i	ii	iii
	(4) endoderm and mesoderm		(2)	iii	ii	iv	i
71.	The contraceptive 'SAHELI'		(3)	<b>i</b> i	iii	i	iv
	(1) is a post-coital contraceptive.		(4)	i	ii	iii	iv
	(2) blocks estrogen receptors in the uterus,	74	3.5	1. (1.	. tacana and		n Column I with those in

preventing eggs from getting implanted.

prevents ovulation in females.

72. The difference between spermiogenesis and

seminiferous tubules.

spermatozoa are formed.

increases the concentration of estrogen and

In spermiogenesis spermatozoa are formed, while in spermiation spermatozoa are released from sertoli cells into the cavity of

In spermiogenesis spermatids are formed, while in spermiation spermatozoa are

In spermiogenesis spermatozoa from sertoli cells are released into the cavity of seminiferous tubules, while in spermiation

In spermiogenesis spermatozoa are formed, while in spermiation spermatids are

is an IUD.

spermiation is

formed.

formed.

belo	w:			
	Colu	mn I		Column II
a.	Glyc	osuria	i.	Accumulation of uric acid in joints
b.	Gout		ii.	Mass of crystallised salts within the kidney
c.	Rena	al calculi	iii.	Inflammation in glomeruli
d.		nerular iritis	iv.	Presence of glucose in urine
	a	b	c	d
(1)	iv	i	ii	iii
(2)	iii	ii	iv	i
(3)	ii	iii	i	iv
(4)	i	ii	iii	iv
Mad	ala the	itoma cri	uon i	n Column I with those in

items given in Column I with those in Column II and select the correct option given below:

	Colu	mnI			Column II
	(Fun	ction)			(Part of Excretory System)
a.	Ultra	afiltrati	on	i.	Henle's loop
b.	Conc of ur	entrati ine	on	ii.	Ureter
c.	Tran	sport o	f	iii.	Urinary bladder
d.	Stor	age of u	rine	iv.	Malpighian corpuscle
				v.	Proximal convoluted tubule
	a	b	c	d	1
(1)	v	iv	i	i	ii
(2)	iv	v	ii	i	ii
(3)	V,	iv	i	i	i i i i i i i i i i i i i i i i i i i
(4)	iv	i	ii	i	ii

75.	Whi	ch of the fo	llowing	gastric cells indirectly	79.		ich of the following events does not occur in				
		in erythropo					gh endoplasmic reticulum ?				
	<b>(1)</b>	Parietal cell	S			(1)	Phospholipid synthesis				
	(2)	Chief cells				(2)	Protein folding Cleavage of signal peptide				
	(3)	Goblet cells				(3)	Protein glycosylation				
	20.0		2000		0.0		12/2 12				
76.	(4) Mucous cells  Match the items given in Column I with those in					Many ribosomes may associate with a smRNA to form multiple copies of a polype					
10.		A CONTRACTOR OF CHARGE TWO	AND AND STREET	e correct option given			ultaneously. Such strings of ribosomes are ned as				
	belo		select the	correct opinion given		(1)	Nucleosome				
	beit	Column I		Column II		(2)	Polysome				
		THE RESERVE OF THE PARTY OF THE	77			(3)	Plastidome				
	a.	Fibrinogen	i.	Osmotic balance		(4)	Polyhedral bodies				
	b.	Globulin	ii.	Blood clotting	81.	Niss	sl bodies are mainly composed of				
	c.	Albumin	iii.	Defence mechanism		(1)	Free ribosomes and RER				
		a b	c			(2)	Proteins and lipids				
	(1)	ii iii	i			(3)	Nucleic acids and SER				
	(2)	iii ii	i			(4)	DNA and RNA				
	(3)	i iii	ii		82.	Whi	ich of these statements is incorrect?				
	(4)	i ii	iii			(1)	Oxidative phosphorylation takes place in outer mitochondrial membrane.				
77.		Which of the following is an occupational respiratory disorder?					Enzymes of TCA cycle are present in mitochondrial matrix.				
	(1)	Emphysema				(3)	Glycolysis operates as long as it is supplied with NAD that can pick up hydrogen atoms				
	(2)	Anthracis				(4)	Glycolysis occurs in cytosol.				
	(3)	Botulism	1 5 1		0.0						
	(4)	Silicosis			83.	(1)					
78.		ium is im raction becau		in skeletal muscle		(2)	chromosomes  Lampbrush – Diplotene bivalents chromosomes				
	(1)	the myosin		tion of bonds between oridges and the actin		(3)	Submetacentric – L-shaped chromososmes chromosomes				
		filament.				(4)	Allosomes – Sex chromosomes				
	(2)	binds to trop active sites of		remove the masking of or myosin.			ch of the following terms describe human tition?				
	(3)	detaches the	e myosir	head from the actin		(1)	Pleurodont, Diphyodont, Heterodont				
		filament.			F. 8	(2)	Thecodont, Diphyodont, Homodont				

(4) activates the myosin ATPase by binding to

(3)

(4)

Pleurodont, Monophyodont, Homodont

Thecodont, Diphyodont, Heterodont

Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing excess water  ich of the following animals does not undergo stamorphosis?  Starfish Earthworm Moth Tunicate  entify the vertebrate group of animals aracterized by crop and gizzard in its digestive stem.  Osteichthyes Amphibia Aves Reptilia  (X/Page 12	97.	(2) (3) (4) Storn (1) (2) (3) (4) The (1) (2) (3) (4) Whi (1) (2) (3) (4) Storn (1) (2) (3) (4) (4) Storn (1) (2) (3) (4) Storn (1) (2) (3) (4)	natal movement is not affected by  CO <sub>2</sub> concentration  Temperature  O <sub>2</sub> concentration  Light  Golgi complex participates in  Activation of amino acid  Fatty acid breakdown  Respiration in bacteria  Formation of secretory vesicles  ch of the following is true for nucleolus?  It is a site for active ribosomal RN synthesis.  Larger nucleoli are present in dividing cells  It takes part in spindle formation.  It is a membrane-bound structure.  stage during which separation of the paire mologous chromosomes begins is  Zygotene  Pachytene  Diakinesis  Diplotene  mata in grass leaf are  Barrel shaped  Dumb-bell shaped  Rectangular  Kidney shaped
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing excess water  ich of the following animals does not undergo stamorphosis?  Starfish Earthworm Moth Tunicate  entify the vertebrate group of animals aracterized by crop and gizzard in its digestive stem.  Osteichthyes Amphibia Aves	95. 96.	(2) (3) (4) Storn (1) (2) (3) (4) The (1) (2) (3) (4) Whii (1) (2) (3) (4) Storn (1) (2) (3) (4) Storn (1) (2) (3)	CO <sub>2</sub> concentration Temperature O <sub>2</sub> concentration Light Golgi complex participates in Activation of amino acid Fatty acid breakdown Respiration in bacteria Formation of secretory vesicles ch of the following is true for nucleolus? It is a site for active ribosomal RN synthesis. Larger nucleoli are present in dividing cells It takes part in spindle formation. It is a membrane-bound structure. stage during which separation of the paire mologous chromosomes begins is Zygotene Pachytene Diakinesis Diplotene mata in grass leaf are Barrel shaped Dumb-bell shaped Rectangular
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing excess water  ich of the following animals does not undergo stamorphosis?  Starfish Earthworm Moth Tunicate  entify the vertebrate group of animals aracterized by crop and gizzard in its digestive estem.  Osteichthyes Amphibia	95. 96.	(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2) (3) (4) Whii (1) The hom (1) (2) (3) (4) Stor (1) (2)	Temperature  O <sub>2</sub> concentration  Temperature  O <sub>2</sub> concentration  Light  Golgi complex participates in  Activation of amino acid  Fatty acid breakdown  Respiration in bacteria  Formation of secretory vesicles  ch of the following is true for nucleolus?  It is a site for active ribosomal RN synthesis.  Larger nucleoli are present in dividing cells  It takes part in spindle formation.  It is a membrane-bound structure.  stage during which separation of the paire cologous chromosomes begins is  Zygotene  Pachytene  Diakinesis  Diplotene  mata in grass leaf are  Barrel shaped  Dumb-bell shaped
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing excess water  ich of the following animals does not undergo etamorphosis?  Starfish Earthworm Moth Tunicate  entify the vertebrate group of animals eracterized by crop and gizzard in its digestive stem. Osteichthyes	95. 96.	(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2) (3) (4) Thi (1) (2) (3) (4) Stor (1) Stor (1)	CO <sub>2</sub> concentration Temperature O <sub>2</sub> concentration Light Golgi complex participates in Activation of amino acid Fatty acid breakdown Respiration in bacteria Formation of secretory vesicles ch of the following is true for nucleolus? It is a site for active ribosomal RN synthesis. Larger nucleoli are present in dividing cells It takes part in spindle formation. It is a membrane-bound structure. stage during which separation of the paire cologous chromosomes begins is Zygotene Pachytene Diakinesis Diplotene mata in grass leaf are Barrel shaped
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing excess water  ich of the following animals does not undergo stamorphosis?  Starfish Earthworm Moth Tunicate  entify the vertebrate group of animals aracterized by crop and gizzard in its digestive stem.	95. 96.	(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2) (3) (4) Whi (1) (2) (3) (4) The hom (1) (2) (3) (4)	CO <sub>2</sub> concentration Temperature O <sub>2</sub> concentration Light Golgi complex participates in Activation of amino acid Fatty acid breakdown Respiration in bacteria Formation of secretory vesicles ch of the following is true for nucleolus? It is a site for active ribosomal RN synthesis. Larger nucleoli are present in dividing cells It takes part in spindle formation. It is a membrane-bound structure. stage during which separation of the paire mologous chromosomes begins is Zygotene Pachytene Diakinesis Diplotene
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing excess water  iich of the following animals does not undergo stamorphosis?  Starfish Earthworm Moth Tunicate  entify the vertebrate group of animals aracterized by crop and gizzard in its digestive	95. 96.	(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2) (3) (4) Whi (1) (2) (3) (4) The hom (1) (2) (3)	CO <sub>2</sub> concentration Temperature O <sub>2</sub> concentration Light Golgi complex participates in Activation of amino acid Fatty acid breakdown Respiration in bacteria Formation of secretory vesicles ch of the following is true for nucleolus? It is a site for active ribosomal RN synthesis. Larger nucleoli are present in dividing cells It takes part in spindle formation. It is a membrane-bound structure. stage during which separation of the paire mologous chromosomes begins is Zygotene Pachytene Diakinesis
Forewings with darker tegmina Presence of caudal styles  sich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  sates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing excess water  sich of the following animals does not undergo etamorphosis?  Starfish Earthworm Moth Tunicate  sentify the vertebrate group of animals	95. 96.	(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2) (3) (4) Whii (1) (2) (3) (4) The hom (1) (2)	CO <sub>2</sub> concentration Temperature O <sub>2</sub> concentration Light Golgi complex participates in Activation of amino acid Fatty acid breakdown Respiration in bacteria Formation of secretory vesicles ch of the following is true for nucleolus? It is a site for active ribosomal RN synthesis. Larger nucleoli are present in dividing cells It takes part in spindle formation. It is a membrane-bound structure. stage during which separation of the paire cologous chromosomes begins is Zygotene Pachytene
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing excess water  ich of the following animals does not undergo stamorphosis?  Starfish Earthworm Moth Tunicate	95. 96.	(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2) (3) (4) Whii (1) The hom (1)	CO <sub>2</sub> concentration Temperature O <sub>2</sub> concentration Light Golgi complex participates in Activation of amino acid Fatty acid breakdown Respiration in bacteria Formation of secretory vesicles ch of the following is true for nucleolus? It is a site for active ribosomal RN synthesis. Larger nucleoli are present in dividing cells It takes part in spindle formation. It is a membrane-bound structure. stage during which separation of the paire cologous chromosomes begins is Zygotene
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing excess water  ich of the following animals does not undergo stamorphosis?  Starfish Earthworm Moth	95.	(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2) (3) (4) Whii (1) (2) (3) (4) The hom	CO <sub>2</sub> concentration Temperature O <sub>2</sub> concentration Light Golgi complex participates in Activation of amino acid Fatty acid breakdown Respiration in bacteria Formation of secretory vesicles ch of the following is true for nucleolus? It is a site for active ribosomal RN synthesis. Larger nucleoli are present in dividing cells It takes part in spindle formation. It is a membrane-bound structure. stage during which separation of the paire
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing excess water  ich of the following animals does not undergo etamorphosis?  Starfish Earthworm	95.	(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2) (3) (4) Whi (1) (2) (3)	CO <sub>2</sub> concentration Temperature O <sub>2</sub> concentration Light Golgi complex participates in Activation of amino acid Fatty acid breakdown Respiration in bacteria Formation of secretory vesicles ch of the following is true for nucleolus? It is a site for active ribosomal RN synthesis. Larger nucleoli are present in dividing cells It takes part in spindle formation. It is a membrane-bound structure. stage during which separation of the paire
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing excess water  ich of the following animals does not undergotamorphosis?  Starfish	95.	(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2) (3) (4) Whi (1) (2) (3) (4)	CO <sub>2</sub> concentration Temperature O <sub>2</sub> concentration Light Golgi complex participates in Activation of amino acid Fatty acid breakdown Respiration in bacteria Formation of secretory vesicles ch of the following is true for nucleolus? It is a site for active ribosomal RN synthesis. Larger nucleoli are present in dividing cells It takes part in spindle formation. It is a membrane-bound structure.
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing excess water  ich of the following animals does not undergo	95. 96.	(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2) (3) (4) Whii (1) (2) (3)	CO <sub>2</sub> concentration Temperature O <sub>2</sub> concentration Light Golgi complex participates in Activation of amino acid Fatty acid breakdown Respiration in bacteria Formation of secretory vesicles ch of the following is true for nucleolus? It is a site for active ribosomal RN synthesis. Larger nucleoli are present in dividing cells It takes part in spindle formation.
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing excess water	95. 96.	(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2) (3) (4) Whi (1) (2)	CO <sub>2</sub> concentration  Temperature O <sub>2</sub> concentration  Light Golgi complex participates in Activation of amino acid Fatty acid breakdown Respiration in bacteria Formation of secretory vesicles ch of the following is true for nucleolus?  It is a site for active ribosomal RN synthesis.  Larger nucleoli are present in dividing cells
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing excess water	95. 96.	(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2) (3) (4) Whii	Temperature  O <sub>2</sub> concentration  Temperature  O <sub>2</sub> concentration  Light  Golgi complex participates in  Activation of amino acid  Fatty acid breakdown  Respiration in bacteria  Formation of secretory vesicles  ch of the following is true for nucleolus?  It is a site for active ribosomal RN
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey having a contractile vacuole for removing	95.	(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2) (3) (4)	CO <sub>2</sub> concentration Temperature O <sub>2</sub> concentration Light Golgi complex participates in Activation of amino acid Fatty acid breakdown Respiration in bacteria Formation of secretory vesicles ch of the following is true for nucleolus?
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion using pseudopodia for capturing prey	95.	(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2) (3)	natal movement is <b>not</b> affected by $\mathrm{CO}_2$ concentration  Temperature $\mathrm{O}_2$ concentration  Light  Golgi complex participates in  Activation of amino acid  Fatty acid breakdown  Respiration in bacteria
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei using flagella for locomotion		(2) (3) (4) Ston (1) (2) (3) (4) The (1) (2)	natal movement is $not$ affected by $CO_2$ concentration  Temperature $O_2$ concentration  Light  Golgi complex participates in  Activation of amino acid  Fatty acid breakdown
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms  iates differ from all other protozoans in having two types of nuclei		(2) (3) (4) Ston (1) (2) (3) (4) The (1)	natal movement is $not$ affected by $CO_2$ concentration  Temperature $O_2$ concentration  Light  Golgi complex participates in  Activation of amino acid
Forewings with darker tegmina Presence of caudal styles ich of the following organisms are known as ef producers in the oceans? Euglenoids Dinoflagellates Cyanobacteria Diatoms iates differ from all other protozoans in		(2) (3) (4) Ston (1) (2) (3) (4) The	natal movement is ${\it not}$ affected by ${\rm CO}_2$ concentration Temperature ${\rm O}_2$ concentration Light Golgi complex participates in
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria Diatoms		(2) (3) (4) Ston (1) (2) (3) (4)	natal movement is $not$ affected by $\mathrm{CO}_2$ concentration Temperature $\mathrm{O}_2$ concentration Light
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates Cyanobacteria	94.	(2) (3) (4) Ston (1) (2) (3)	natal movement is $not$ affected by $\mathrm{CO}_2$ concentration Temperature $\mathrm{O}_2$ concentration
Forewings with darker tegmina Presence of caudal styles  ich of the following organisms are known as ef producers in the oceans?  Euglenoids Dinoflagellates	94.	(2) (3) (4) Ston (1) (2)	natal movement is ${\it not}$ affected by ${\rm CO}_2$ concentration ${\rm Temperature}$
Forewings with darker tegmina Presence of caudal styles ich of the following organisms are known as ef producers in the oceans? Euglenoids	94.	(2) (3) (4) Ston (1)	natal movement is $not$ affected by $\mathrm{CO}_2$ concentration
Forewings with darker tegmina Presence of caudal styles ich of the following organisms are known as ef producers in the oceans?	94.	(2) (3) (4) Ston	natal movement is not affected by
Forewings with darker tegmina Presence of caudal styles ich of the following organisms are known as	94.	(2) (3) (4)	
Forewings with darker tegmina Presence of caudal styles		(2) (3)	NADH
Forewings with darker tegmina		(2)	
TO SUBSTITUTE STATE OF THE STAT			NADPH
TO SUBSTITUTE STATE OF THE STAT		(1)	Oxygen ATP
9 <sup>th</sup> abdominal segment			etion of photosynthesis?
Presence of a boat shaped sternum on the	93.		ch of the following is not a product of ligh
Presence of anal cerci		(4)	Mycobacterium
nale cockroach from a female cockroach?		(3)	Nostoc
ich of the following features is used to identify			Saccharomyces
Chelone	92.		ch among the following is <b>not</b> a prokaryote?  Oscillatoria
	00	XX71	
		(4)	
Macropus		(3) (4)	carbonyl and methyl
		(2) (3) (4)	carbonyl and phosphate carbonyl and methyl
Psittacula		(3)	carbonyl and methyl
	Camelus  Chelone  citch of the following features is used to identify	nich of the following features is used to identify	cich of the following features is used to identify  (1) (2)

	(5)				9 120 L		
99.	Polle	n grains can be stored for several years in	106.	Selection (1)	t the <i>correct</i> match: Francois Jacob and	100	Lac operon
		d nitrogen having a temperature of		(1)	Jacques Monod		Date of the second
	(1)	− 160°C		(0)	Alec Jeffreys	-	Streptococcus
	(2)	-120°C		(2)	Alec Jenreys		p <mark>neumoniae</mark>
	(3)	−196°C		(3)	Matthew Meselson		Pisum sativum
	(4)	−80°C		(3)	and F. Stahl	1363	
100.	Oxyg	gen is <b>not</b> produced during photosynthesis by		10	Alfred Hershey and	- '	TMV
	(1)	CONTRACT CON		(4)	Martha Chase		1111
	(2)	Green sulphur bacteria	102	F01		for	comiconcervative
	(3)	Cycas	107.	The	experimental proof cation of DNA was fir	et sho	wn in a
	(4)	Nostoc		(1)	Virus	50 5110	
101.	Doul	ole fertilization is		(2)	Fungus		
	(1)	Syngamy and triple fusion		(3)	Plant		
	(2)	Fusion of two male gametes of a pollen tube		(4)	Bacterium		
	N-0	with two different eggs	100	1.500	ct the <i>correct</i> statem	ent :	
	(3)	Fusion of two male gametes with one egg	108.		Transduction was di		ed by S Altman
	(4)	Fusion of one male gamete with two polar		(1)			
		nuclei		(2)			
102.	Whi	ch one of the following plants shows a very		(3)	Spliceosomes take p		
	close	e relationship with a species of moth, where		(4)	Punnett square was	devel	loped by a British
	none	e of the two can complete its life cycle without			scientist.		5 12/15
	the	other?	109.		ch of the following	g pa	irs is wrongly
	(1)	Viola	141	mat	ched?		200
	(2)	Hydrilla	KH	(1)	T.H. Morgan		
	(3)	Banana		(2)	Starch synthesis in		
	(4)	Yucca		(3)	XO type sex		Grasshopper
103.	Whi	ch of the following elements is responsible for			determination		
	mai	ntaining turgor in cells?		(4)	ABO blood grouping	:	: Co-dominance
	(1)	Calcium	110.	Offs	sets are produced by		
	(2)	Magnesium		(1)	Parthenogenesis		
	(3)	Potassium	P	(2)	Meiotic divisions		
	(4)	Sodium		(3)	Parthenocarpy		
104.	In v	which of the following forms is iron absorbed		(4)	Mitotic divisions		
	by I	plants?	111.	Wh	ich of the following	flower	s only once in it
	(1)	Both ferric and ferrous			-time ?		
	(2)	Ferric		(1)	Papaya		
	(3)	Free element		(2)	Bamboo species		
	(4)	Ferrous		(3)	Mango		
105		at is the role of NAD+ in cellular		(4)	Jackfruit		
105			112.	Wh	ich of the following	has	proved helpful in
			4	pre	serving pollen as foss	ls?	
	(1)			(1)	Sporopollenin		
	(2)			(2)	Pollenkitt		
			teit	(3)	Oil content		
			1	(4)	Cellulosic intine		
105.		at is the role of NAD <sup>+</sup> in cellular piration?  It is the final electron acceptor for anaerobic respiration.  It functions as an enzyme.  It is a nucleotide source for ATP synthesis.  It functions as an electron carrier.	112.	Wh pre (1) (2) (3)	ich of the following serving pollen as foss Sporopollenin Pollenkitt Oil content	has ils?	proved helpful

- and organisations without authorisation from the concerned country and its people is called
  - Bioexploitation
  - (2) Bio-infringement
  - Biodegradation (3)
  - Biopiracy (4)
- 114. Which of the following is commonly used as a vector for introducing a DNA fragment in human lymphocytes?
  - pBR 322
  - Retrovirus (2)
  - λ phage
  - Ti plasmid
- 115. The correct order of steps in Polymerase Chain Reaction (PCR) is
  - Denaturation, Annealing, Extension
  - Extension, Denaturation, Annealing
  - Denaturation, Extension, Annealing
  - Annealing, Extension, Denaturation
- 116. A 'new' variety of rice was patented by a foreign company, though such varieties have been present in India for a long time. This is related to
  - (1) Basmati
  - (2) Co-667
  - Lerma Rojo (3)
  - (4) Sharbati Sonora
- 117. Select the correct match:
  - (1) G. Mendel
- Transformation
- Ribozyme (2)
- Nucleic acid
- T.H. Morgan
- Transduction
- (4) F<sub>2</sub> × Recessive parent Dihybrid cross
- 118. In India, the organisation responsible for assessing the safety of introducing genetically modified organisms for public use is
  - Genetic Engineering Appraisal Committee (GEAC)
  - Indian Council of Medical Research (ICMR)
  - Research Committee Genetic Manipulation (RCGM)
  - and Industrial Council for Scientific Research (CSIR)

- 113. Use of bioresources by multinational companies 119. Which of the following is a secondary pollutant?
  - 0,
  - CO
  - SO2 (3)
  - (4) CO<sub>2</sub>
  - 120. Natality refers to
    - Number of individuals entering a habitat
    - Death rate
    - Number of individuals leaving the habitat
    - Birth rate
  - 121. Niche is
    - the functional role played by the organism where it lives
    - all the biological factors in the organism's environment
    - the range of temperature that the organism (3) needs to live
    - the physical space where an organism lives
  - 122. World Ozone Day is celebrated on
    - 22<sup>nd</sup> April
    - 5<sup>th</sup> June
    - 16<sup>th</sup> September
    - 21st April
  - 123. In stratosphere, which of the following elements acts as a catalyst in degradation of ozone and release of molecular oxygen?
    - Oxygen
    - Carbon
    - Fe
    - Cl (4)
  - What type of ecological pyramid would be obtained with the following data?

Secondary consumer: 120 g

Primary consumer: 60 g

Primary producer: 10 g


- Upright pyramid of biomass
- Inverted pyramid of biomass
- Upright pyramid of numbers
- Pyramid of energy

5. Pneumatophores occur in	Match the items given in Column I with those in Column II and select the <i>correct</i> option given
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	below: Column I Column II
(2) Halophytes	Continue 1
(3) Carnivorous plants	A. Herbarium i. It is a place having a
(4) Free-floating hydrophytes	collection of preserved
26. Sweet potato is a modified	plants and animals.
(1) Rhizome	b. Key ii. A list that enumerates
A second	methodically all the
	species found in an area
	with brief description
(4) Adventitious root	aiding identification.
27. Secondary xylem and phloem in dicot stem are	c. Museum iii. Is a place where dried and
produced by	pressed plant specimens
(1) Axillary meristems	mounted on sheets are
(2) Apical meristems	kept.
(3) Phellogen	d. Catalogue iv. A booklet containing a list
(4) Vascular cambium	of characters and their
rn: 1 of the following statements is correct?	alternates which are
argually inprancticu in	helpful in identification of
Coope and Cedrus.	various taxa.
at anglosed by ovary wall in	a b c d
(2) Ovules are not enclosed by gymnosperms.	i i
il and grimposperms.	(1) 6
a de la la hoterosporous, while Salvilla	(2) 1 1 1 1
is homosporous.	(3) 11 17 111
	(4) 111 11
129. Casparian strips occur in	133. After karyogamy followed by meiosis, spores an
(1)	produced exogenously in
(2) Epidermis	- I
(3) Cortex	The state of the s
(4) Pericycle	
statement:	
(1) Mitochondria are the powerhouse of the cell	3.77
in all kingdoms except Monera.	134. Which one is wrongly matched?
il to progent in members of rungi	(1) Unicellular organism - Chilorette
1 Diantag	(2) Uniflagellate gametes - Polysiphonia
ti- and locomotory and feeding	(3) Gemma cups - Marchanta
tmietures in Sporozoalis.	(4) Biflagellate zoospores - Brown algae
halong to Basidiomycetes.	
(4) Mushrooms belong to bush growth are	135. Winged pollen grains are present in
131. Plants having little or no secondary growth are	(1) Pinus
(1) Cycads	(2) Mustard
(2) Grasses	(3) Mango
(3) Conifers	(4) Cycas
Desiduous angiosperms	ROUGH WORK En

- pipe is equal to the third harmonic of a closed organ pipe. If the length of the closed organ pipe is 20 cm, the length of the open organ pipe is
  - 16 cm (1)
  - 13.2 cm (2)
  - 12.5 cm
  - 8 cm (4)
- 137. At what temperature will the rms speed of oxygen molecules become just sufficient for escaping from the Earth's atmosphere?

Mass of oxygen molecule (m) =  $2.76 \times 10^{-26}$  kg Boltzmann's constant  $k_B = 1.38 \times 10^{-23} \ J \ K^{-1})$ 


- $1.254 \times 10^4 \text{ K}$
- $2.508 \times 10^{4} \text{ K}$
- $5.016 \times 10^4 \text{ K}$ (3)
- (4)  $8.360 \times 10^4 \text{ K}$
- 138. The efficiency of an ideal heat engine working between the freezing point and boiling point of water, is
  - 12.5% (1)
  - 26.8% (2)
  - 6.25% (3)
  - (4)
- 139. The volume (V) of a monatomic gas varies with its temperature (T), as shown in the graph. The ratio of work done by the gas, to the heat absorbed by it, when it undergoes a change from state A to state B, is



- (1)
- (2)
- (3)

- 136. The fundamental frequency in an open organ 140. A metallic rod of mass per unit length 0.5 kg m<sup>-1</sup> is lying horizontally on a smooth inclined plane which makes an angle of 30° with the horizontal. The rod is not allowed to slide down by flowing a current through it when a magnetic field of induction 0.25 T is acting on it in the vertical direction. The current flowing in the rod to keep it stationary is
  - 11.32 A
  - 7·14 A (2)
  - 14.76 A
  - 5.98 A (4)
  - 141. An inductor 20 mH, a capacitor 100 µF and a resistor 50 Ω are connected in series across a source of emf, V = 10 sin 314 t. The power loss in the circuit is
    - 1.13 W (1)
    - 0.79 W
    - 2.74 W (3)
    - 0.43 W (4)
  - 142. A thin diamagnetic rod is placed vertically between the poles of an electromagnet. When the current in the electromagnet is switched on, then the diamagnetic rod is pushed up, out of the horizontal magnetic field. Hence the rod gains gravitational potential energy. The work required to do this comes from
    - (1) the induced electric field due to the changing magnetic field
    - the current source (2)
    - the lattice structure of the material of the
    - the magnetic field
  - 143. Current sensitivity of a moving coil galvanometer is 5 div/mA and its voltage sensitivity (angular deflection per unit voltage applied) is 20 div/V. The resistance of the galvanometer is
    - 500 Ω (1)
    - 40 Ω
    - $250 \Omega$
    - $25 \Omega$


- 144. A moving block having mass m, collides with another stationary block having mass 4m. The lighter block comes to rest after collision. When the initial velocity of the lighter block is v, then the value of coefficient of restitution (e) will be
  - (1) 0.4
  - (2) 0.5
  - (3) 0.8
  - (4) 0.25
- 145. A body initially at rest and sliding along a frictionless track from a height h (as shown in the figure) just completes a vertical circle of diameter AB = D. The height h is equal to




- (1)  $\frac{5}{4}$  D
- (2)  $\frac{3}{2}$  D
- $(3) \quad \frac{7}{5} \, D$
- (4) D
- 146. Which one of the following statements is incorrect?
  - Coefficient of sliding friction has dimensions of length.
  - (2) Rolling friction is smaller than sliding friction.
  - (3) Frictional force opposes the relative motion.
  - (4) Limiting value of static friction is directly proportional to normal reaction.
- 147. Three objects, A: (a solid sphere), B: (a thin circular disk) and C: (a circular ring), each have the same mass M and radius R. They all spin with the same angular speed ω about their own symmetry axes. The amounts of work (W) required to bring them to rest, would satisfy the relation
  - $(1) \quad W_{A} > W_{C} > W_{B}$
  - $(2) \quad W_C > W_B > W_A$
  - $(3) \quad W_{B} > W_{A} > W_{C}$
  - $(4) \quad W_A > W_B > W_C$

- 148. Unpolarised light is incident from air on a plane surface of a material of refractive index '\mu'. At a particular angle of incidence 'i', it is found that the reflected and refracted rays are perpendicular to each other. Which of the following options is correct for this situation?
  - (1)  $i = tan^{-1} \left(\frac{1}{\mu}\right)$
  - (2) Reflected light is polarised with its electric vector parallel to the plane of incidence
  - $(3) \quad i = \sin^{-1} \left(\frac{1}{u}\right)$
  - (4) Reflected light is polarised with its electric vector perpendicular to the plane of incidence
  - 149. In Young's double slit experiment the separation d between the slits is 2 mm, the wavelength λ of the light used is 5896 Å and distance D between the screen and slits is 100 cm. It is found that the angular width of the fringes is 0·20°. To increase the fringe angular width to 0·21° (with same λ and D) the separation between the slits needs to be changed to
    - (1) 1·7 mm
    - (2) 1.8 mm
    - (3) 2·1 mm
    - (4) 1.9 mm
    - 150. An astronomical refracting telescope will have large angular magnification and high angular resolution, when it has an objective lens of
      - (1) small focal length and small diameter
      - (2) small focal length and large diameter
      - (3) large focal length and large diameter
      - (4) large focal length and small diameter

151. In the circuit shown in the figure, the input voltage  $V_i$  is 20 V,  $V_{BE}$  = 0 and  $V_{CE}$  = 0. The values of  $I_B$ ,  $I_C$  and  $\beta$  are given by





- (1)  $I_B = 40 \mu A$ ,  $I_C = 5 mA$ ,  $\beta = 125$
- (2)  $I_B = 40 \mu A$ ,  $I_C = 10 \text{ mA}$ ,  $\beta = 250$
- (3)  $I_B = 20 \mu A$ ,  $I_C = 5 mA$ ,  $\beta = 250$
- (4)  $I_B = 25 \mu A$ ,  $I_C = 5 mA$ ,  $\beta = 200$
- 152. In a p-n junction diode, change in temperature due to heating
  - (1) affects the overall V I characteristics of p-n junction
  - (2) affects only reverse resistance
  - (3) does not affect resistance of p-n junction
  - (4) affects only forward resistance
- 153. In the combination of the following gates the output Y can be written in terms of inputs A and B as





- (1) A + B
- (2) A.B
- (3)  $\overline{A.B} + A.B$
- (4)  $A \cdot \overline{B} + \overline{A} \cdot B$

- 154. A carbon resistor of  $(47 \pm 4.7)$  k $\Omega$  is to be marked with rings of different colours for its identification. The colour code sequence will be
  - (1) Green Orange Violet Gold
  - (2) Violet Yellow Orange Silver
  - (3) Yellow Green Violet Gold
  - (4) Yellow Violet Orange Silver
- 155. A set of 'n' equal resistors, of value 'R' each, are connected in series to a battery of emf 'E' and internal resistance 'R'. The current drawn is I. Now, the 'n' resistors are connected in parallel to the same battery. Then the current drawn from battery becomes 10 I. The value of 'n' is
  - (1) 9
  - (2) 10
  - (3) 20
  - (4) 11
- 156. A battery consists of a variable number 'n' of identical cells (having internal resistance 'r' each) which are connected in series. The terminals of the battery are short-circuited and the current I is measured. Which of the graphs shows the correct relationship between I and n?









- have the same volume. The first wire has cross-sectional area A and the second wire has cross-sectional area 3A. If the length of the first wire is increased by  $\Delta l$  on applying a force F, how much force is needed to stretch the second wire by the same amount?
  - (1) F
  - (2) 9 F
  - (3) 4 F
  - (4) 6 F
- 158 A sample of 0.1 g of water at 100°C and normal pressure (1.013 × 10<sup>5</sup> Nm<sup>-2</sup>) requires 54 cal of heat energy to convert to steam at 100°C. If the volume of the steam produced is 167.1 cc, the change in internal energy of the sample, is
  - (1) 84·5 J
  - (2)104·3 J
  - (3) 42.2 J
  - 208·7 J
- 159 A small sphere of radius & falls from rest in a viscous liquid. As a result, heat is produced due to viscous force. The rate of production of heat when the sphere attains its terminal velocity, is proportional to
  - (1)  $r^4$

  - (3)
  - (4)
- 160% The power radiated by a black body is P and it radiates maximum energy at wavelength, λ<sub>0</sub>. If the temperature of the black body is now changed so that it radiates maximum energy at wavelength  $\frac{3}{4}\lambda_0$ , the power radiated by it 164. becomes nP. The value of n is
  - 81 256

  - 256 81


- 157 Two wires are made of the same material and 161. An electron falls from rest through a vertical distance h in a uniform and vertically upward directed electric field E. The direction of electric field is now reversed, keeping its magnitude the same. A proton is allowed to fall from rest in it through the same vertical distance h. The time of fall of the electron, in comparison to the time of fall of the proton is
  - equal. (1)
  - smaller -(2)
  - 10 times greater (3)
  - 5 times greater
  - A pendulum is hung from the roof of a sufficiently high building and is moving freely to and fro like a simple harmonic oscillator. The acceleration of the bob of the pendulum is 20 m/s2 at a distance of 5 m from the mean position. The time period of oscillation is
    - 1 s
    - $2\pi s$ (2)
    - 28
  - A tuning fork is used to produce resonance in a glass tube. The length of the air column in this tube can be adjusted by a variable piston. At room temperature of 27°C two successive resonances are produced at 20 cm and 73 cm of column length. If the frequency of the tuning fork is 320 Hz, the velocity of sound in air at 27°C is
    - 300 m/s (1)
    - 330 m/s (2)
    - (3)350 m/s
    - 339 m/s (4)
  - The electrostatic force between the metal plates of an isolated parallel plate capacitor C having a charge Q and area A, is
    - inversely proportional to the distance between the plates.
    - independent of the distance between the (2)plates.
    - proportional to the square root of the distance between the plates.
    - linearly proportional to the distance between the plates.

Sux

- threshold frequency), is incident on a metal plate, the maximum velocity of electrons emitted is v1. When the frequency of the incident radiation is increased to  $5v_0$ , the maximum velocity of electrons emitted from the same plate is v2. The ratio of v1 to v2 is
  - (1) 2:1
  - 1:2
  - 4:1 (3)
  - (4) 1:4
- 166. For a radioactive material, half-life 10 minutes. If initially there are 600 number of nuclei, the time taken (in minutes) for the disintegration of 450 nuclei is
  - 15 (1)
  - 20 (2)
  - 30 (3)
  - (4) 10
- 167. The ratio of kinetic energy to the total energy of an electron in a Bohr orbit of the hydrogen atom,

  - 1:-2 (1) (2)1:1
  - (3) 2:-1
  - (4) 1:-1
- 168. An electron of mass m with an initial velocity  $\overrightarrow{V} = V_0 \overrightarrow{i} (V_0 > 0)$  enters an electric field  $\overrightarrow{E} = -E_0 \hat{i}$  ( $E_0 = \text{constant} > 0$ ) at t = 0. If  $\lambda_0$  is its de-Broglie wavelength initially, then its de-Broglie wavelength at time t is
  - (1)  $\lambda_0$


- 165. When the light of frequency  $2v_0$  (where  $v_0$  is 169. A solid sphere is in rolling motion. In rolling energy (Kt) as well as rotational kinetic energy  $(K_r)$  simultaneously. The ratio  $K_t$ ;  $(K_t + K_r)$  for the sphere is
  - (1) 2;5
  - (2)7:10
  - 10:7 (3)
  - 5:7
  - 170. If the mass of the Sun were ten times smaller and the universal gravitational constant were ten times larger in magnitude, which of the following is not correct?
    - 'g' on the Earth will not change.
    - Raindrops will fall faster.
    - Time period of a simple pendulum on the Earth would decrease.
    - Walking on the ground would become more difficult.
  - 171. A solid sphere is rotating freely about its symmetry axis in free space. The radius of the sphere is increased keeping its mass same. Which of the following physical quantities would remain constant for the sphere?
    - Angular momentum (1)
    - Angular velocity (2)
    - Rotational kinetic energy (3)
    - Moment of inertia
  - 172. The kinetic energies of a planet in an elliptical orbit about the Sun, at positions A, B and C are KA, KB and KC, respectively. AC is the major axis and SB is perpendicular to AC at the position of the Sun S as shown in the figure. Then



- (1)  $K_B > K_A > K_C$
- (2)  $K_A < K_B < K_C$
- (3)  $K_B < K_A < K_C$
- $K_A > K_B > K_C$

- 1736 The refractive index of the material of a prism is  $\sqrt{2}$  and the angle of the prism is 30°. One of the two refracting surfaces of the prism is made a mirror inwards, by silver coating. A beam of monochromatic light entering the prism from the other face will retrace its path (after reflection from the silvered surface) if its angle of incidence on the prism is
  - (1) zero
  - (2) 60°
  - (3) 30°
  - (4) 45°
- 174 The magnetic potential energy stored in a certain inductor is 25 mJ, when the current in the inductor is 60 mA. This inductor is of inductance
  - (1) 13·89 H
  - (2) 0·138 H
  - (3) 1·389 H
  - (4) 138·88 H
- An object is placed at a distance of 40 cm from a concave mirror of focal length 15 cm. If the object is displaced through a distance of 20 cm towards the mirror, the displacement of the image will be
  - (1) 36 cm towards the mirror
  - (2) 30 cm away from the mirror
  - (3) 30 cm towards the mirror
  - (4) 36 cm away from the mirror
- 176. An em wave is propagating in a medium with a velocity \( \bar{V} = \bar{V} \bar{i} \). The instantaneous oscillating electric field of this em wave is along +y axis. Then the direction of oscillating magnetic field of the em wave will be along
  - (1) -x direction
  - (2) z direction
  - (3) y direction
  - (4) + z direction

177. A block of mass m is placed on a smooth inclined wedge ABC of inclination θ as shown in the figure. The wedge is given an acceleration 'a' towards the right. The relation between a and θ for the block to remain stationary on the wedge



- (1)  $a = g \tan \theta$
- (2)  $a = \frac{g}{\cos e c \theta}$
- (3)  $a = g \cos \theta$
- (4)  $a = \frac{g}{\sin \theta}$
- 178. A student measured the diameter of a small steel ball using a screw gauge of least count 0.001 cm. The main scale reading is 5 mm and zero of circular scale division coincides with 25 divisions above the reference level. If screw gauge has a zero error of -0.004 cm, the correct diameter of the ball is
  - (1) 0.529 cm
  - (2) 0.521 cm
  - (3) 0.053 cm
  - (4) 0.525 cm
- 179. The moment of the force,  $\vec{F} = 4\hat{i} + 5\hat{j} 6\hat{k}$  at (2, 0, -3), about the point (2, -2, -2), is given by
  - (1)  $-7\hat{i} 4\hat{j} 8\hat{k}$
  - (2)  $-8\hat{i} 4\hat{j} 7\hat{k}$
  - (3)  $-7\hat{i} 8\hat{j} 4\hat{k}$
  - (4)  $-4\hat{i} \hat{j} 8\hat{k}$
- 180. A toy car with charge q moves on a frictionless horizontal plane surface under the influence of a uniform electric field E. Due to the force qE, its velocity increases from 0 to 6 m/s in one second duration. At that instant the direction of the field is reversed. The car continues to move for two more seconds under the influence of this field. The average velocity and the average speed of the toy car between 0 to 3 seconds are respectively
  - (1) 1.5 m/s, 3 m/s
  - (2) 2 m/s, 4 m/s
  - (3) 1 m/s, 3·5 m/s
  - (4) 1 m/s, 3 m/s

Sux